

Information topic	Details
Indicator name	<b>Days with extreme rainfall / Days with soil moisture deficit</b>
Domain and topic	Climate change: Extreme rainfall and drought
Indicator definition and units	<p><b>Extreme rainfall days</b> are defined as days with rainfall amounts exceeding the 95<sup>th</sup> percentile, which was calculated from daily rainfall amounts across the climate normal period from 1991-2020.</p> <p><b>Annual rainfall anomaly</b> is defined as the difference between the quantity of rainfall received in a given year, and the average annual rainfall across the climate normal period from 1991-2020.</p> <p><b>Number of days with soil moisture deficit 'dry days'</b>: A day is 'severely' dry if the deficit of moisture in the soil is between 110-130mm. A day is 'extremely' dry if the deficit is equal to or greater than 130mm.</p>
Data source	- NIWA Datahub: <a href="https://data.niwa.co.nz/">https://data.niwa.co.nz/</a>
Numerator	<ul style="list-style-type: none"> <li>• Annual number of days with extreme rainfall by Territorial Authority (TA)</li> <li>• Annual number of days with a soil moisture deficit by Territorial Authority (TA)</li> </ul>
Methodology	<p>Climate stations were selected based on their proximity to the population-weighted centroid for a TA as well as completeness of data for the period 1991-2023. One weather station per TA was selected.</p> <p>Using the population-weighted centroid coordinates for each TA, we looked at weather stations within a 25km radius. The weather station closest to the centroid was selected, provided it was currently operating and had a long record of data (i.e., minimum of 10 years of data). Where there was insufficiently complete data or the station was closed, we then examined the next closest weather station, and so on until the 'best fit' was found.</p> <p><i>For rainfall:</i></p> <p>In three cases, a climate station is used for two TAs (Hamilton/Waikato, Lower Hutt/Porirua, Tauranga/Western Bay of Plenty).</p> <p><i>For soil moisture deficit:</i></p> <p>In four cases, a climate station is used for two TAs (Hamilton/Waikato, Lower Hutt/Porirua, New Plymouth/Stratford, Tauranga/Western Bay of Plenty).</p> <p><i>For both indicators</i></p> <p>In two cases, the only suitable climate station was currently closed</p>

	<p>and an exception was made (Kaipara, Opotiki).</p> <p>The population-weighted centroid of a TA was calculated from 2018 Census data, using the geographic centroid of statistical area 1 (SA1, small Census area description) weighted by their usual resident population. The selection of stations for this indicator will be reviewed once population data from the 2023 Census becomes available.</p> <p>The most recent Climate Normal for New Zealand was calculated as an average over the 30-year period 1991-2020 (all available data from all TAs was included). This average number acted as a benchmark against which current or recent observations were compared (i.e., anomalies).</p> <p><i>Adjustments for disrupted data collection</i></p> <p>If a climate station's data had over 25% missing data in 2023, it was excluded from the analysis. In publications <b>before and after 2023</b>, the cutoff was set at 10%. The wider margin applies only to data for 2023 (i.e. to the Surveillance Report published in 2024), to allow for disruption to data collection due to Cyclone Gabrielle.</p>
Time period and time scale	<p>Annual; from 1991 onwards</p> <p>Three-year moving averages, from 2000–2002 onwards</p>
Spatial Coverage	National and by TA
Measures of frequency	<ul style="list-style-type: none"> <li>- Average number of days per year with extreme rainfall and soil moisture deficit</li> <li>- Number of days per year with extreme rainfall and soil moisture deficit, by TA</li> <li>- 1991-2020 baseline average number of days with extreme rainfall or soil moisture deficit</li> <li>- Annual rainfall anomaly, by TA</li> </ul>
Limitations of indicator	<ul style="list-style-type: none"> <li>- There will be geographic variation across a TA that is not represented for these indicators because we have used one weather station per TA.</li> </ul>
Limitations of data source	<p>Some of the selected weather stations have missing data, usually due to starting collection after the year 1991.</p> <p>Several TAs do not have a suitably representative weather station, or lack data to report, particularly for Soil Moisture Deficit in the South Island.</p>
Created by	Environmental Health Intelligence New Zealand, Massey University
Related indicators	<ul style="list-style-type: none"> <li>- Number of days below 0°C</li> <li>- Number of days over 25°C</li> <li>- Notifications of salmonellosis, cryptosporidiosis, campylobacteriosis and giardiasis</li> </ul>

For more information	<ul style="list-style-type: none"> <li>- <a href="https://www.niwa.co.nz/climate/nz-drought-monitor/droughtindicatormaps">https://www.niwa.co.nz/climate/nz-drought-monitor/droughtindicatormaps</a></li> <li>- Ministry for the Environment &amp; Stats NZ. 2020. New Zealand's Environmental Reporting Series: Our atmosphere and climate 2020. Wellington: Ministry for the Environment &amp; Stats NZ.</li> </ul>
References	