

Number and density of livestock in **New Zealand**

This factsheet presents indicators of the number and density of livestock (sheep, dairy cattle, beef cattle and deer) in New Zealand. Livestock plays an important role in New Zealand's economy, though poorly managed livestock production can significantly impact water quality/recreational water quality.



Overall, livestock numbers declined by more than a quarter between 2002 and 2022.



Dairy cattle were the only livestock type whose numbers increased between 2002 and 2022, although they appear to be declining in recent years.



Though sheep numbers decreased by one-third between 2002–2022, sheep still outnumbered all other livestock types combined by a factor of just over two to one as of 2022.



The Manawatū-Whanganui region had the greatest density of livestock, with 424.4 animals per km² of farmland in 2022.

Agricultural activity has a major effect on recreational water

Agricultural use of land can cause bodies of water that are used by people for recreational activities (such as swimming, boating, and fishing) to become contaminated in several ways (Ministry for the Environment & Statistics New Zealand 2023):


- Storm run-off of livestock waste from farms into water sources can affect water quality by introducing pathogens such as *Campylobacter*, *Cryptosporidium*, *E.coli*, and *Giardia* contained in sheep and cow dung. These pathogens can cause gastrointestinal or respiratory diseases, especially in immunocompromised people (Devane et al 2018).
- Biological contaminants such as pesticides and pharmaceuticals used to treat livestock have been detected in waterways internationally and are shown to potentially increase development and spread of antimicrobial resistance (Alderton et al 2021).
- Excess nutrients, such as nitrogen from fertiliser or livestock urine, can be washed into waterways by rain and reduce water quality by promoting the growth of potentially toxic algae.
- Irrigation for intensive farming of livestock is one of the largest uses of freshwater in New Zealand, which may lower the water levels in rivers. Lowered water levels can interrupt river flow and prevent harmful contaminants from being 'flushed' out of a waterway.

Additionally, agriculture is a notable contributor to greenhouse gas emissions. The methane (CH₄) emissions make up over two thirds of New Zealand's agricultural emissions, making it one of the largest sources of emissions in the country (Agricultural Greenhouse Gas Research Centre 2023).

Total livestock numbers continue to decrease

There were 36.2 million stock animals in New Zealand in 2022. Between 2002 and 2022, the number of stock animals in New Zealand decreased by about a third, from 50.9 million to 36.2 million (Figure 1, Table 1). The bulk of the decline relates to decreasing sheep numbers, which reduced by just over a third (36.0%) during this period. Despite the reduction in numbers, sheep still outnumbered all other livestock types combined by a factor of just over two to one as of 2022.

The number of beef cattle decreased by 13.2% over the 20-year period between 2002 and 2022, but has been gradually increasing again since 2015. Dairy cattle numbers increased by 18.9% overall, but the numbers peaked in 2014 and have been slowly decreasing. The majority of the increase occurred in the South Island, where the number of dairy cattle almost doubled. Deer numbers declined markedly, with a 51.8% decrease between 2002 and 2022.



Source: Statistics New Zealand 2023. The data for 2002, 2007, 2012, 2017, and 2022 were collected from Agricultural Production Census data, whereas in between years were collected from Agricultural Production Statistics surveys.

Livestock type	2002 (millions)	2021 (millions)	2022 (millions)	Percent change (2002–2022)	Percent change (2021-2022)
Sheep	39.57	25.73	25.33	-36.0%	-1.6%
Dairy	5.16	6.19	6.14	+18.9%	-0.8%
Beef	4.49	3.97	3.90	-13.2%	-1.7%
Deer	1.65	0.82	0.79	-51.8%	-2.5%
Total	50.87	36.70	36.16	-28.9%	-1.5%

Source: Statistics New Zealand 2023

Total livestock density

The overall livestock density in New Zealand in 2022 was 274.2 animals per km² of farmland. However, livestock density varies by region. In 2022, the Manawatū-Whanganui region had the highest density of livestock (424.4 animals per km²). The density of livestock types also varied between regions, reflective of variations in farming practices between regions.







Data for this indicator

Data comes from Statistics New Zealand's Agricultural Production Statistics, which contain the results of the Agricultural Production Censuses and Agricultural Production Surveys conducted from the year 2002 onwards.

For descriptive information about the data, see the **Metadata Sheet.**

References

Alderton, I., Palmer, B. R., Heinemann, J. A., Pattis, I., Weaver, L., Gutiérrez-Ginés, M. J., ... & Tremblay, L. A. (2021). The role of emerging organic contaminants in the development of antimicrobial resistance. *Emerging Contaminants*, 7, 160-171.

Devane, M. L., Weaver, L., Singh, S. K., & Gilpin, B. J. (2018). Fecal source tracking methods to elucidate critical sources of pathogens and contaminant microbial transport through New Zealand agricultural watersheds–A review. *Journal of Environmental Management*, 222, 293-303.

Agricultural Greenhouse Gas Research Centre. 2023. *The science of methane*. https://www.nzagrc.org.nz/domestic/methane-research-programme/the-science-of-methane/ (accessed Oct 2023).

Ministry for the Environment (MfE) & Statistics New Zealand. 2015. *New Zealand's Environmental Reporting Series: Environment Actearoa 2015.*Available from www.mfe.govt.nz and www.mfe.govt.nz (accessed July 2017).

Statistics New Zealand. 2023a. Agricultural Production Census. Data available from from www.stats.govt.nz/information-releases/ (accessed May 2023)

Statistics New Zealand. 2023b. *Agricultural Production Statistics: June 2022 (final)*. Data available from www.stats.govt.nz/information-releases/agricultural-production-statistics-year-to-june-2022-final (accessed May 2023).

Previous factsheet:

2021

Other related topics include:

Water-borne diseases related to recreational water

Agricultural land use

<u>Leptospirosis notifications</u>

Disclaimer

Drought and rainfall

Environmental Health Intelligence NZ – Rapu Mātauranga Hauora mo te Taiao - Aotearoa, makes no warranty, express or implied, nor assumes any legal liability or responsibility for the accuracy, correctness, completeness or use of any information that is available on this factsheet

Author

To get in touch with the author <u> ehinz@massey.ac.nz</u>

Citation

Environmental Health Intelligence NZ, 2023. *Livestock numbers and density.* Wellington: Environmental Health Intelligence NZ, Massey University.

Q <u>Visit our website</u>

Subscribe to our newsletter

🛩 f in