

Sudden unexpected death in infancy (SUDI)

This report presents information on rates of sudden unexpected death in infancy (SUDI) from 2010 to 2019 (the most recent data available) in New Zealand.

Key facts

- 46 babies died from SUDI in 2019, up from 37 deaths the previous year. There has been no improvement in SUDI rates since 2012.
- Pacific and Māori babies had five times the rate of SUDI as European/Other babies in 2015–19.
- Babies of younger mothers (younger than 25 years) had higher SUDI rates than babies born to mothers in older age groups.
- The SUDI rate for babies living in the most socioeconomically deprived areas (NZdep 2013 quintile 5) was more than seven times as high as babies in the least deprived areas (quintile 1).

Maternal smoking doubles the risk of SUDI


Exposure to [second-hand smoke](#) has been shown to increase the risk of SUDI in infants (under one year old) (US Department of Health and Human Services 2007, Anderson and Cook 1997). Having a mother who smokes also doubles their risk of dying from SUDI (Zhang and Wang 2013; Anderson and Cook 1997). The estimated number of SUDI deaths attributable to maternal smoking decreased from 7 in 2010 to 3 in 2019 (based on Mason and Borman (2016) methodology). Similarly, the latest data shows that [maternal smoking at two weeks postnatal](#) fell from 13.7% in 2009 to 7.3% in 2021.

In 2002–2010, New Zealand had a high SUDI rate compared with other developed countries (Taylor et al 2015).

46 babies died from SUDI in 2019

In 2019, 46 babies less than one year of age (<1 year) (0.8 per 1,000 live births; 95%CI 0.6–1.0) died from SUDI, up from 37 the previous year (Figure 1). In total, 451 babies died from SUDI between 2010 and 2019.


Figure 1: Number of SUDI deaths in children aged <1 year, 2010–2019



Source: Health New Zealand – Te Whatu Ora, 2023

The SUDI rate declined from 2010 to 2012, followed by a slight increase until 2014 (Figure 2). Since then, it has remained relatively steady, with no significant improvement. In 2017, the New Zealand government launched the National SUDI prevention programme. A target was set to reduce the SUDI rate from 0.7 per 1,000 live births to 0.1 per 1,000 live births by 2025 (Ministry of Health 2017).

Figure 2: SUDI deaths in children aged <1 year, 2010–2019 (rate per 1,000 live births)




Note: 95% confidence intervals have been presented as vertical bars.

Source: Health New Zealand – Te Whatu Ora, 2023

Pacific and Māori babies dying at a higher rate

Of the 46 babies that died from SUDI in 2019, 29 were Māori (63%), 8 were Pacific (17%), 8 were European (17%), and 1 was Asian (2%). In 2015–19, Pacific (1.5 per 1,000 live births; 95%CI 1.1–2.0) and Māori babies (1.4 per 1,000 live births; 95%CI 1.2–1.7) had five times the rate of SUDI as European/Other babies (0.3 per 1,000 live births; 95%CI 0.2–0.4) (Figure 3).

Figure 3: SUDI deaths in children aged <1 year, by ethnic group (prioritised), 2015–19 (rate per 1,000 live births)




Note: 95% confidence intervals have been presented as vertical bars.
Source: Health New Zealand – Te Whatu Ora, 2023

Babies of younger mothers have higher rates of SUDI

In 2015-19, SUDI rates were higher for babies whose mothers were younger than 25 years old (Figure 4).

Figure 4: SUDI deaths in children aged <1 year, by maternal age, 2015–19 (rate per 1,000 live births)




Note: 95% confidence intervals have been presented as vertical bars.
Source: Health New Zealand – Te Whatu Ora, 2023

Higher SUDI rates in the most socioeconomically deprived areas

In 2015–19, the SUDI rate for babies living in the most deprived areas (NZDep2013 quintile 5) was statistically significantly higher (1.5 per 1,000 live births; 95%CI 1.2–1.8) than those living in other quintiles. Babies living in the most deprived areas had more than seven times the rate of SUDI as babies living in the least deprived areas (quintile 1).

Figure 5: SUDI deaths in children aged <1 year, by NZDep2013 quintile, 2015–19 (rate per 1,000 live births)




Note: 95% confidence intervals have been presented as vertical bars.
Source: Health New Zealand – Te Whatu Ora, 2023

Tairāwhiti and Whanganui districts had high SUDI rates in 2015–19

In 2015–19, Tairāwhiti (1.6 per 1,000 live births; 95% CI 0.6–3.6) and Whanganui (1.6 per 1,000 live births; 95% CI 0.7–3.4) districts had high SUDI rates (Figure 6).

Figure 6: SUDI deaths in children aged <1 year, by district, 2015–19 (rate per 1,000 live births)



Note: 95% confidence intervals have been presented as vertical bars. *The rate is suppressed for Lakes, Hutt Valley, Wairarapa, Nelson Marlborough, West Coast, and South Canterbury districts due to counts less than five.
Source: Health New Zealand – Te Whatu Ora, 2023

Data for this indicator

This indicator includes the most recent data available from the Fetal and Infant Deaths web tool published by Health New Zealand – Te Whatu Ora in December 2023.

The indicator presents data related to sudden unexpected death in infancy (SUDI), defined as deaths in children aged less than one year old, with an underlying cause of death in the following ICD-10AM codes:

- R95 – sudden infant death syndrome (SIDS)
- R96 – other sudden death, cause unknown
- R98 – unattended death
- R99 – other ill-defined and unspecified causes
- W75 – accidental suffocation and strangulation in bed
- W78 – inhalation of gastric contents
- W79 – inhalation and ingestion of food causing obstruction of respiratory tract

This definition follows the recommendations of the Child and Youth Mortality Review Committee (2009).

Mortality rates are presented as deaths per 1,000 live births.

Calculating the attributable SUDI deaths due to maternal smoking

The estimated number of SUDI deaths linked to maternal smoking is based on the methodology used by Mason and Borman (2016). For this report, we have used more recent SUDI and maternal smoking data, along with an updated relative risk of 1.97 from Zhang and Wang (2013). Mason and Borman's 2016 study, by contrast, used a relative risk of

1.94 from Anderson and Cook (1997). Despite these changes, the differences in the estimates are minimal.

For additional information, see the [Metadata](#) sheet.

References

Anderson HR, Cook D. 1997. Passive smoking and sudden infant death syndrome: review of the epidemiological evidence. *Thorax* 52: 1003–09.

Child and Youth Mortality Review Committee, Te Rōpū Arotake Auau Mate o te Hunga Tamariki, Taiohi. 2009. *Fifth Report to the Minister of Health: Reporting mortality 2002–2008*. Wellington: Child and Youth Mortality Review Committee.

Health New Zealand – Te Whatu Ora. 2023. *Fetal and Infant Deaths web tool*. URL: <https://tewhatuora.shinyapps.io/fetal-and-infant-deaths-web-tool/> (accessed 7 May 2024).

Mason K, Borman B. 2016. Burden of disease from second-hand smoke exposure in New Zealand. *New Zealand Medical Journal*. 129 (1432): 16–25.

Ministry of Health. 2017. *National SUDI prevention programme launched*. 2017. [Press release]. URL: <https://www.beehive.govt.nz/release/national-sudi-prevention-programme-launched> (accessed 6 July 2021).

Taylor BJ, Garstang J, Engelberts A et al. 2015. International comparison of sudden unexpected death in infancy rates using a newly proposed set of cause-of-death codes. *Archives of Disease in Childhood* 100(11): 1018–23. DOI: 10.1136/archdischild-2015-308239.

US Department of Health and Human Services. 2007. *Children and Secondhand Smoke Exposure. Excerpts from The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health.

Zhang K, Wang X. 2013. Maternal smoking and increased risk of sudden infant death syndrome: A meta-analysis. *Legal Medicine* 15: 115–121.

Explore geographic data on interactive dashboards:

[Indoor Environment domain dashboard](#)

[EHINZ dashboard](#)

Previous surveillance reports:

[2022](#)

[2021](#)

[2019](#)

Other related topics include:

[Maternal smoking at two weeks postnatal](#)

[Second-hand smoke exposure](#)

[Health burden due to second-hand smoke exposure](#)

[Household crowding](#)

[Home heating](#)

Disclaimer

Environmental Health Intelligence NZ – Rapu Mātauranga Hauora mo te Taiao - Aotearoa, makes no warranty, express or implied, nor assumes any legal liability or responsibility for the accuracy, correctness, completeness or use of any information that is available in this surveillance report.

Author

The author of this report is Helene Marsters, ehinz@massey.ac.nz

Citation

Environmental Health Intelligence. 2024. *Sudden unexpected death in infancy (SUDI)* [Surveillance Report]. Wellington: Environmental Health Intelligence NZ, Massey University.

[Visit the EHINZ website](#)

[Subscribe to our newsletter](#)